overall, but is not enough to reverse the declining population trend without additional effort (Pabian et al. 2015).

Prior research indicates that farm raised ring-necked pheasants use habitat with ample cover in all seasons. Fall habitat use varies among regions, however, farm-raised birds generally favor idle, untilled cropland (Smith et al. 1999, Gatti et al. 1989; Leif 2005), shrub lands and wetlands (Klinger and Riegner 2008, Marco et al. 2010). In its ring-necked pheasant management plan, the Pennsylvania Game Commission also cites plantings of cool season grasses and forbs as a preferred fall environment (Klinger and Riegner 2008). In addition, from research studies done by Neilson et al. (2008), they found that ring-necked pheasant population size was a good indicator of conservation reserve success along agricultural areas in the United States.

The purpose of this project was to determine both the habitat usage preference and the home range extent of male and female pen-raised pheasants upon reintroduction to a wild landscape using radio-telemetry and GIS data analysis. As demonstrated by Marcstrom et al. (1989) and Matthews et al. (2012), fitting pheasants with necklace transmitters and tracking via radio telemetry provides an innocuous, and efficient method to generate the necessary data for spatial analysis. Our null hypothesis was that both male and female pheasants would exhibit similar landscape usage preferences and range extents during the project period.

**MATERIA-(1.9 T ID 3ErNETEs A9ErNETED 3TD 3HOrNETEe)MCI.ces i19an**

into Lancaster County Central Park (LCCP), mid-afternoon on September 18th, and were equally spaced along a 750 m transect (Figure 2). The pheasant release transect traveled the margin of an open, contiguous, shrubby, 13-hectare early successional field. The field was mostly surrounded on all sides by tree canopy. Following release, the pheasants were tracked using a hand-held antenna and radio-receiver manufactured by Telonics Inc. (http://telonics.com/). Once pheasant locations were confirmed in the field, their positions were marked using hand-held Garmin eTrex Venture GPS units (www.garmin.com). Pheasant positions were marked at least twice daily, Morning (6:30 – 9:30 am), Noon (11:00 am – 2:00 pm), Evening (2:00 – 5:00 pm) and/or Dusk (5:00 – 7:00 pm).

Data was collected from September through October 2015, or until pheasant mortality. Spatial data was analyzed using the Biotas® software program (http://www.ecostats.com/web/Biotas). Home range estimations were made using the 95% fixed-kernel home range estimator with least squares cross validation. Land cover raster satellite data was obtained from the Pennsylvania Geospatial Data Clearinghouse (www.pasda.psu.edu) for Lancaster County. Land cover data was divided into four categories: grass/shrub, paved, tree canopy and water. The field defined as “grass/shrub”, comprised 51% of the project area. The deciduous forest was defined as “tree canopy”, and comprised 43% of the project area. “Paved” surface area covered 3% of the project area. The final category “water” accounted for a nearby river (Conestoga River) and occupied 3% of the project area (Figure 2). Land cover selection (i.e., habitat selection) analysis was conducted for pen-raised ring-necked pheasants using Goodness-of-fit comparisons of expected number of pheasant locations in each habitat or land cover type based on habitat availability. This was compared to observed pheasant locations in each habitat type represented by 95% confidence intervals (Neu et al. 1974). This analysis was

conducted using the Biotas® software program to calculate the $X^2$

## DISCUSSION

The average range extent of pheasants in this project (6.04 ha) was markedly lower than demonstrated in other similar studies. Ramey et al. (2006) found that pen-raised birds averaged a home range size of 117.2 ha, while Smith et al. (1999) showed that home range size can range from 35 – 150 ha. Considerations such as population density, regional climate, land usage and/or sampling method contribute to the variability in range size (Smith et al. 1999). Since home range size increased with survival duration during this project (Table 1), it is reasonable to infer that the surviving birds' home range would continue to expand a

NEUTEL, W., BEALEY, C.E. 1993. Effects of land use on breeding pheasant density. Journal of Applied Ecology. Vol. 30 (3): 465-477

PABIAN, S. E., A. M. WILSON, S. R. KLINGER, and M. C. BRITTINGHAM, 2015. Pennsylvania's Conservation Reserve Enhancement Program benefits ring-necked pheasants but not enough to reverse declines. Journal of Wildlife Management 79(4): 641 – 646.

RAMEY, C. A., J. B. BOURASSA, and M. S. FURUTA. 2006. Survival and Movement Ecology of Ring-Necked Pheasants in Northern California Agricultural Areas. Proc. 22nd

**Table 1. Home range size estimates based on a fixed kernel estimator with least squares cross validation and survival duration of six pen-raised ring-necked pheasants released and radio-monitored in Lancaster County Park from September through October, 2015.**

| Pheasant ID | Sex | Home Range (hectares) | Survival (days) | Usage Rate (hectares/day) |
|---|---|---|---|---|
| 15 | Female | 5.48 | 40** | 0.137 |
| 25 | Female | 1.65 | 12 | 0.138 |
| 35 | Female | 3.37 | 10 | 0.337 |
| 45 | Male | | | |

**FIGURES**

Figure 1. Male ring-necked pheasant fitted with a 12-gram Advanced Telemetry Solutions (https://atstrack.com/) necklace radio transmitter.




Figure 3. Analysis of habitat usage preference (Grass/Shrub, Paved, Tree Canopy, Water) of six pen-raised ring-necked pheasants (two male, four female) released into Lancaster County Central Park in September 2015. Squares indicate the proportion of habitat available on the project area and the 95% confidence intervals indicate the proportion of habitat used by pheasants.




Figure 5. Analysis of habitat usage preference (Grass/Shrub, Paved, Tree Canopy, Water) of two male pen-raised ring-necked pheasants released into Lancaster County Central Park in September 2015. Squares indicate the proportion of habitat available on the project area and the 95% confidence intervals indicate the proportion of habitat used by pheasants.